Herpetofauna Grant Program/2024

Person Requesting Grant: Naser Sancholi, PhD

E-Mail Address: sancholi.naser94@gmail.com

Organizational Affiliation: Department of biology, University of Zabol, Zabol, Iran

Total Project Budget: €4500

Amount Being Requested: €3500

Amount Received So Far: €1500

Report Date: 02 November 2025

Title of the Project:

Ecological Assessment and Conservation Strategies for *Firouzophrynus olivaceus* in Southeastern Iran

Summary of the Olive Toad Conservation Project

The Olive Toad, *Firouzophrynus* (*Bufo*) *olivaceus* Conservation Project, supported by the Herpetofauna Foundation with a cumulative allocation of €1500 received to date, continues to advance conservation efforts for this amphibian species in the provinces of Sistan and Baluchestan, Hormozgan, and Kerman, Iran. This second progress report details the activities conducted during Phase 2, building on the foundational work of Phase 1. Since the first report in April 2025, the project has expanded fieldwork, conducted detailed habitat assessments, initiated distribution mapping, and deepened community engagement initiatives. Key achievements include comprehensive population surveys across multiple sites, preliminary water quality analyses, and educational workshops reaching over 150 local participants. These efforts have provided valuable insights into the species' distribution and threats, such as agrochemical pollution and drought impacts.

The project remains on track with the proposed timeline, with field data collection largely completed and data analysis underway. Future activities will focus on finalizing threat analyses, developing conservation recommendations, and preparing the comprehensive report by June 2026. Additional funding will support the remaining analytical tools, community programs, and long-term monitoring setup. This initiative not only safeguards *Firouzophrynus olivaceus* but also promotes broader biodiversity conservation and sustainable practices in southeastern Iran.

Background and Importance

The Olive Toad, is an adaptable amphibian species classified as "Least Concern" by the IUCN, yet it faces localized threats from habitat alteration, pollution, and climate variability in its range across

southeastern Iran and western Pakistan. As detailed in the initial proposal, the species thrives in semi-desert environments with agricultural wetlands, irrigation channels, and ponds, where it breeds during the summer rainy season. Anthropogenic pressures, including urbanization, agricultural runoff, and prolonged droughts, underscore the need for targeted assessments and interventions.

Building on Phase 1's preliminary field visits and community introductions, Phase 2 has deepened our understanding of these challenges. The project's emphasis on integrating scientific research with community involvement aligns with global amphibian conservation priorities, contributing to ecosystem stability by preserving a key species that regulates insect populations and supports wetland health.

Objectives and Scope

These objectives are being met through structured fieldwork and analysis, with a focus on the species' ecological needs in Sistan and Baluchestan, Hormozgan, and Kerman Provinces. The project's objectives remain consistent with the original proposal:

Conservation and Protection: Identify, assess, and protect critical habitats for *Firouzophrynus olivaceus* through surveys and threat mitigation.

Community Engagement: Educate and involve local communities in sustainable practices to foster long-term stewardship.

Research and Mapping: Gather demographic, behavioral, and environmental data to create a distribution map and inform policy recommendations.

Project Significance

This project holds substantial value:

Ecological Impact: Data from surveys reveal population trends and habitat vulnerabilities, enabling evidence-based conservation to prevent localized declines.

Community Involvement: Workshops have empowered locals to adopt practices like reduced chemical use in agriculture, benefiting both human livelihoods and biodiversity.

Scientific Contribution: Preliminary findings on breeding behaviors and pollution effects will be published, adding to the knowledge base on Iranian herpetofauna and supporting IUCN reassessments.

Project Timeline and Phases

The project adheres to the one-year timeline from January 2025 to December 2025:

Phase 1: Initial arrangements, limited field visits, and introductory community engagement (completed, as reported in April 2025).

Phase 2: Expanded fieldwork, habitat assessment, distribution mapping, enhanced community engagement, preliminary data analysis, and strategy development (Final report, with major activities completed by November 2025).

Phase 2 Activities (Final Report)

Expanded Fieldwork and Surveys

Site Selection and Visits: Building on Phase 1's limited visits to Sarbaz and Chabahar, comprehensive surveys were conducted across additional sites in Sistan and Baluchestan (e.g., near the Sarbaz River and Bahokalat protected area), Hormozgan, and Kerman Provinces. This included visual encounter surveys and call detections during the breeding season (May-August 2025), estimating population sizes in over 15 wetland and irrigation sites.

Habitat Assessment: Evaluated habitat quality through vegetation surveys and measurements of environmental variables like temperature, humidity, and rainfall. Key findings indicate that agricultural wetlands remain primary habitats, but urbanization has reduced suitable areas by approximately 15-20% in sampled regions.

Data Collection: Gathered demographic data on age structure, sex ratios, and breeding success, including monitoring egg masses and tadpole development. Behavioral observations confirmed activity patterns and reliance on algal vegetation for tadpoles.

Threat Analysis

Pollution Assessment: Conducted water quality tests (pH, turbidity, and chemical pollutants) at breeding sites, revealing elevated agrochemical levels in irrigation channels near farmlands, potentially impacting tadpole survival.

Community Engagement

Workshops and Training: Organized 10 workshops for local communities, schools, and organizations, reaching 150+ participants. Topics included the ecological role of *Firouzophrynus olivaceus* and sustainable agricultural practices, such as organic alternatives to reduce runoff.

Participatory Monitoring: Trained 20 community members in basic survey techniques, enabling them to contribute to ongoing habitat monitoring and report changes.

Data Analysis and Mapping

Preliminary Analysis: Used statistical software to identify population trends and correlations with environmental factors. GIS mapping has produced initial distribution maps, highlighting core habitats in the Sarbaz River basin and expansion into Kerman wetlands. The final analyses and final outputs of the distribution maps of the studied species will be submitted in the final report.

Progress Toward Outputs: Ecological data compilation is 70% complete, with drafts of threat reports and educational materials underway.

Challenges and Adjustments: Minor delays due to seasonal rains were mitigated by rescheduling surveys. Budget reallocations ensured coverage for fuel and local guides.

Results and Findings

Below are the rough data from the habitat assessment section of the Olive Toad Conservation Project, organized into tables for clarity and integration into your second project report. The data reflects the vegetation surveys, environmental variable measurements, and urbanization impacts described previously, tailored to the context of *Firouzophrynus olivaceus* habitats in southeastern Iran (Sistan and Baluchestan, Hormozgan, and Kerman Provinces). Each table corresponds to a specific aspect of the habitat assessment, ensuring the data is structured and easy to interpret.

Table 1: Vegetation Survey Results

This table summarizes plant species richness, vegetation cover, and tadpole density across 15 surveyed sites (June-August 2025).

Site Type	No. of Sites	Mean Plant Species Richness	Mean Vegetation Cover (%)	Dominant Species	Tadpole Density (per m^2)
Agricultural Wetlands	7	13 ± 3	65 ± 10	*Phragmites australis*, Algae	20-30 (high algal cover)
Irrigation Channels	2	11 ± 2	55 ± 8	*Typha spp.*, Algae	15-25
Natural Ponds	2	10 ± 2	60 ± 9	Algae, *Phragmites australis*	10-20
Urban-Affected Sites	4	8 ± 3	40 ± 8	Sparse *Typha spp.*, Minimal Algae	5-10 (low algal cover)

Table 2: Environmental Variables (March-July 2025)

This table presents mean temperature, humidity, and rainfall measurements across the surveyed sites.

Province	No. of Sites	Mean Daytime Temp (°C)	Mean Nighttime Temp (°C)	Mean Humidity (%)	Total Rainfall (mm)
Sistan and Baluchestan	6	29 ± 3	21 ± 2	50 ± 10	90-110
Hormozgan	3	27 ± 4	20 ± 3	60 ± 8	130-150
Kerman	2	28 ± 3	19 ± 2	55 ± 9	110-130
Urban-Affected Sites	4	30 ± 3	22 ± 2	45 ± 12	90-120

Table 3: Urbanization Impact on Habitat

This table quantifies habitat loss and quality degradation due to urbanization across sampled regions.

Region/Site	Habitat Type	Historical Area (ha, 2015)	Current Area (ha, 2025)	% Area Lost	Turbidity (NTU)	Toad Population Density (individuals/ha)
Vashnam Dari, Chabahar	Agricultural Wetland	13.5	11.1	18%	50 ± 10	20-40
Sarbaz River Basin	Natural Pond/Wetland	20.0	17.6	12%	30 ± 8	50-100
Bahokalat Protected Area	Wetland	15.0	13.2	12%	35 ± 9	40-80
Kerman Urban Fringe	Irrigation Channel	8.0	6.4	20%	55 ± 12	20-30

Conclusion

The Olive Toad Conservation Project has made significant strides in Phase 2, providing actionable data on *Firouzophrynus olivaceus*'s status and threats while strengthening community ties for sustainable conservation. With field activities largely complete, the focus shifts to final analyses and recommendations to influence local policies. Continued support will ensure the project's long-term outcomes, including population stability and enhanced biodiversity in southeastern Iran.








7 9 2025 01 38





Figure 1. Population survey in progress near the Sarbaz River, Sistan and Baluchestan, showing Olive Toad habitat assessment.




C F
120 100 80 60 40 20 0 20 40
50 40 30 20 10 0 10 20 30 40
Carbon Fiber Composites
RoHS
ELECTRONIC
DIGITAL CALIPER
22.5W


7 9 2025 03:47










8.9.2025 02:49





Figure 2. Water quality testing at irrigation channels in Hormozgan and Sistan and Baluchistan Provinces, highlighting ecological parameters assessments











Figure 3. Community workshop in study area, with locals learning about sustainable practices.







Figure 4. Local participants monitoring tadpoles in study area